

and clinical symptoms being responsive to L-dopa and appearing stable across time. However, the general concept of disease progression is probably too heterogeneous to assume that a 5-year period of 10 patients using our clinical and TMS paradigms is sufficient to allow firm conclusions. Therefore, a follow-up multimodal examination is planned in another 5 years. ●

Acknowledgments: This study was supported by the Else Kröner-Fresenius Foundation (2018_A55) and the German Research Foundation (WE 5919/2-1, SFB936 project C5, and FOR2488). Support by the European Reference Network–Rare Neurological Diseases (Project 739510) is also gratefully acknowledged.

Data Availability Statement

The data that support the findings of this study are available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.

Anne Weissbach, MD,^{1,2*} Annika Steinmeier, MS,¹
 Martje G. Pauly, MD,^{1,2,3} Duha M. Al-Shorafat, MD,^{4,5}
 Gerard Saranza, MD,^{4,6} Anthony E. Lang, MD,⁴
 Norbert Brüggemann, MD,³ Vera Tadic, MD,³
 Christine Klein, MD,² Katja Lohmann, PhD,²
 Matt J.N. Brown, PhD,⁷ Christian Beste, PhD,⁸
 Alexander Münchau, MD,¹ and Tobias Bäumer, MD¹

¹Institute of Systems Motor Science, University of Lübeck, Lübeck, Germany, ²Institute of Neurogenetics, University of Lübeck, Lübeck, Germany, ³Department of Neurology, University Hospital Schleswig Holstein, Lübeck, Germany, ⁴Edmond J. Safra Program in Parkinson's Disease and the Morton and Gloria Shulman Movement Disorders Clinic, Toronto Western Hospital, Toronto, Ontario, Canada, ⁵Neuroscience Department, Jordan University of Science and Technology, Irbid, Jordan, ⁶Section of Neurology, Department of Medicine, Chong Hua Hospital, Cebu City, Philippines, ⁷Department of Kinesiology, California State University Sacramento, Sacramento, California, USA, and ⁸Cognitive Neurophysiology, Department of Child and Adolescent Psychiatry, Faculty of Medicine of the TU Dresden, Dresden, Germany

References

1. Ichinose H, Ohye T, Takahashi E, et al. Hereditary progressive dystonia with marked diurnal fluctuation caused by mutations in the GTP cyclohydrolase I gene. *Nat Genet* 1994;8(3):236–242.
2. Hanajima R, Nomura Y, Segawa M, Ugawa Y. Intracortical inhibition of the motor cortex in Segawa disease (DYT5). *Neurology* 2007; 68(13):1039–1044.
3. Huang YZ, Trender-Gerhard I, Edwards MJ, Mir P, Rothwell JC, Bhatia KP. Motor system inhibition in dopa-responsive dystonia and its modulation by treatment. *Neurology* 2006;66(7):1088–1090.
4. Weissbach A, Baumer T, Brüggemann N, et al. Premotor-motor excitability is altered in dopa-responsive dystonia. *Mov Disord* 2015;30:1705–1709.

Supporting Data

Additional Supporting Information may be found in the online version of this article at the publisher's web-site.

Focal Subthalamic Atrophy after Long-Term Deep Brain Stimulation in Parkinson's Disease

Deep brain stimulation (DBS) is a surgical procedure that is increasingly being applied to selected patients with movement disorders, and other conditions. In Parkinson's disease (PD), DBS is particularly effective against disabling and pharmacotherapy-resistant motor symptoms.¹ DBS has been effectively applied to the nucleus ventralis intermedialis of the thalamus, the subthalamic nucleus (STN),² or the internal globus pallidus.^{3,4} We previously reported that DBS induces no significant brain damage between 3 and 70 months (nearly 3 years; mean 33 months) of electrode implantation, with the exception of a mild chronic tissue reaction around the lead track.⁴

Here we present postmortem neuropathological findings of four additional patients who underwent long-term DBS for a period of up to 16 years (96–192 months) and compared them with nine additional PD patients that did not undergo DBS. Neurosurgical procedures and neuropathological workup are detailed in Supplementary file S1 and results are summarized in Table S1.

In all cases, we observed a mild to moderate gliofibrillar capsule around the track of the electrode, mild surrounding gliosis with occasional distortion of axonal profiles (Fig. 1B1–B4), reactive microglia and occasional (hemosiderin) pigment. In patient 3 we found several foreign body multinucleated giant cells (Fig. 1C2, arrow) around the tip of the electrode. In all cases, the tip of the electrode was located within or beside the STN (Fig. 1). In two patients (patient 1, patient 4) there was a distortion of the medial part of the nucleus with moderate shrinkage of its volume (Fig. 1A1–A4). Compared to the lateral part of the STN and to the STN of nine non-DBS treated PD patients, there was a moderate astroglia, microglial

© 2021 The Authors. *Movement Disorders* published by Wiley Periodicals LLC on behalf of International Parkinson and Movement Disorder Society

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Key Words: Parkinson's disease; deep brain stimulation; subthalamic nucleus

***Correspondence to:** Prof. François Alesch, MD, Department of Neurosurgery, Medical University of Vienna, Waehringer Guertel 18-20, 1090 Vienna, Austria; E-mail: francois.alesch@meduniwien.ac.at
 Prof. Christine Haberler, MD, Division of Neuropathology and Neurochemistry, Department of Neurology, Medical University of Vienna, Waehringer Guertel 18-20, 1090 Vienna, Austria; E-mail: christine.haberler@meduniwien.ac.at

Financial Disclosure/Conflict of Interests

The authors report no conflict of interest. No specific funding was received for the present study.

Received: 30 April 2021; **Accepted:** 2 May 2021

Published online 12 June 2021 in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/mds.28653




FIG. 1. Histological features at the subthalamic nucleus (STN) and track (for more detailed features see Figure S2). **(A1–A4)** STN (encircled) of patient 1 with an adjacent hole caused by the tip of the electrode, seen at the upper part of the figure (asterisk). The medial part of the STN (upper arrow left) shows oedematous change with small vacuoles (**A1**, **A2**: Klüver Barrera [KB] stain, the blue colour represents myelin sheaths). **A3**: There is a reduction of neuronal density in MAP2 immunohistochemistry with smaller residual neurons and condensed neuronal processes on the left side. **A4**: This medial area of the STN also shows an increase in microglia and a moderate chronic fibrillary astrocytic gliosis (**A4**). **(B1–B4)** In all cases, a mild tissue reaction was observed around the lead track with formation of a gliofibrillar capsule already visible on hematoxylin eosin (HE)-stained sections (patient 1 as representative for all cases) with reduction of myelin sheaths (**B2**, Klüver-Barrera), increase of reactive, amoeboid shaped microglia (**B3**) and mild-to-moderate chronic fibrillar gliosis (**B4**). **(C1–C4)** In patient 3 there was also an accumulation of multinucleated giant cells at the tip of the electrode (**C1** box and **C2** in higher magnification) associated with a proliferation of Rosenthal fibres around the hole (**C3**). There was also more prominent immunoreaction for human leukocyte antigen-DR isotype (HLA-DR) (**C4**). **(D1–D4)** Abundant alpha-synuclein immunoreactive Lewy bodies and Lewy neurites were detected in several brainstem nuclei of all cases (**D1** substantia nigra/patient 1; **D2** dorsal motor nucleus of the vagal nerve/patient 2; **D3** locus coeruleus/patient 3; and **D4** dorsal motor nucleus of the vagal nerve/patient 4) confirming the diagnosis of Parkinson's disease. **(d1–d4)**: Mild-to-moderate amount of alpha-synuclein immunoreactive cell processes around the track/tip of the electrode (**d1** to **d4** correspond to patients 1 to 4, respectively). Original magnifications: A1, A2, A3, A4, B1, B2, B3, B4, C1, C4: x40; D1, D2, D3, D4, d1–d4: x100; C2, C3, D1, d4: x200. [Color figure can be viewed at wileyonlinelibrary.com]

activation, coarsening of neuronal processes, and a change of neuronal shape, size, and density (Fig. S2). These changes were consistent with a moderate degeneration, particularly of the medial aspect – limbic part – of the STN. This was also observed to a milder degree in patient 2, while in patient 3 only tiny and gliotic fragments of the STN were identified. These changes did not seem to be related to the implantation site, the disease duration, or the intensity of stimulation.

In summary, our observations show that in contrast to short-duration DBS, chronic DBS may elicit moderate

degeneration of the stimulated target neuronal tissue, here the STN, after up to 16 years of stimulation. It remains unclear whether these changes are related to chronic depolarization block or due to an alteration of the synaptic conduction or other causes including the mere physical proximity of a foreign body, and is a matter of speculation. However, there are descriptions of positive effects of DBS on synaptic function and neurogenesis.^{5,6} In contrast, along the lead track, we found no major changes besides a gliofibrotic capsule, as described previously,^{4,7} and occasional foreign body reaction

at the tip of the electrode. Mild to moderate alpha-synuclein aggregates were also identified within neuronal processes in the immediate surroundings of the canal and the tip of the electrode. Although not very prominent, these findings might suggest that chronic noxae may increase alpha-synuclein aggregation.⁸

Our findings are important particularly because the application of DBS for different diseases is being increasingly performed on younger patients.¹ DBS is, however, a stable and effective long-term therapy. ■

Acknowledgments: We are indebted to brain donors and families for the generous donation of brain tissue for diagnostic and research purposes. We thank the pathologists for coordinating autopsies and the treating neurologists of the patients presented here: Prim. Dr. Ekkehard Spuller, Institut für klinische Pathologie und Molekularpathologie LKH Villach; Prof. Dr. Martin Klimpfner, Pathologisch-Bakteriologisches Institut Kaiser-Franz-Josef Spital; Prim. Dr. Dieter Volc, Neurologische Abteilung der Confraternität; and Prim. Dr. Christa Freibauer, Institut für klinische Pathologie und Molekularpathologie, LK Mistelbach, all in Austria. We thank Daniel C. Bradley for proofreading the English version of the manuscript.

Ellen Gelpi, MD,¹  Christine Haberler, MD,^{1*} 
 Alexander Micko, MD,²  Andrea Polt, BSc,¹
 Andreas Amon, MScD,² Karl Rössler, MD,²  and
 François Alesch, MD^{2*}

¹*Division of Neuropathology and Neurochemistry, Department of Neurology, Medical University of Vienna, Vienna, Austria, and*

²*Department of Neurosurgery, Medical University of Vienna, Vienna, Austria*

Data Availability Statement

The data that supports the findings of this study are available in the supplementary material of this article.

References

- Schuepbach WM, Rau J, Knudsen K, Volkmann J, et al. Neurostimulation for Parkinson's disease with early motor complications. *N Engl J Med* 2013;368:610–622.
- Hamel W, Köppen JA, Alesch F, Antonini A. Targeting of the subthalamic nucleus for deep brain stimulation: a survey among Parkinson disease specialists. *World Neurosurg* 2017;99:41–46.
- Timmermann L, Jain R, Chen L, Maarouf M, et al. Multiple-source current steering in subthalamic nucleus deep brain stimulation for Parkinson's disease (the VANTAGE study): a non-randomised, prospective, multicentre, open-label study. *Lancet Neurol* 2015;14:693–701.
- Haberler C, Alesch F, Mazal PR, Pilz P, et al. No tissue damage by chronic deep brain stimulation in Parkinson's disease. *Ann Neurol* 2000;48:372–376.
- Pohodich AE, Yalamanchili H, Raman AT, Wan YW, et al. Forniceal deep brain stimulation induces gene expression and splicing changes that promote neurogenesis and plasticity. *Elife* 2018;7:7.
- Hammond C, Ammari R, Bioulac B, Garcia L. Latest view on the mechanism of action of deep brain stimulation. *Mov Disord* 2008;23:2111–2121.
- Reddy GD, Lozano AM. Postmortem studies of deep brain stimulation for Parkinson's disease: a systematic review of the literature. *Cell Tissue Res* 2018;373:287–295.
- Pal GD, Ouyang B, Serrano G, Shill HA, et al. Arizona study of aging neurodegenerative disorders. Comparison of neuropathology in Parkinson's disease subjects with and without deep brain stimulation. *Mov Disord* 2017;32:274–277.

Supporting Data

Additional Supporting Information may be found in the online version of this article at the publisher's web-site.